

Predicting the invasive potential of the cladoceran *Daphnia lumholtzi* Sars, 1885 (Crustacea: Cladocera: Daphniidae) in the Neotropics: Are generalists threatened and relicts protected by their life-history traits?

Sousa F., Palaoro A., Elmoor-Loureiro L., Kotov A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017, Page Press Publications All rights reserved. Invasive species are one of the major threats to biodiversity, which is aggravated in poorly known groups, such as cladocerans. *Daphnia lumholtzi* Sars (Cladocera: Anomopoda: Daphniidae) is currently invading the Neotropical region, and there are few records of this process. Our goal was to predict the invasive scenario for *D. lumholtzi* in the Neotropics using species distribution modelling and to assess the climatic overlap of the invader with the native species. We trained our MaxEnt model using occurrence records from native and invaded areas and projected it in the Neotropics. Additionally, we compared the climatic niche of some native species with the invader's niche. Our model showed high environmental suitability in areas connected by the lowland Paraná River Basin (southwestern Brazil, eastern Argentina and Uruguay), in south-central Chile and Atlantic coastal areas. Widely distributed native species showed climatic overlap with the invader, while relict species did not. *Daphnia lumholtzi* thrives in warm and stable environments (e.g. the Paraná River basin), which of concern because the invader could already be spreading in that area. Native species could suffer due to climatic niche similarity, while natural barriers and local environmental conditions may protect relict species. We urge the need for further studies to understand this invasion process more fully.

<http://dx.doi.org/10.4081/jlimnol.2016.1571>

Keywords

Biodiversity conservation, Dispersion, Invasive species, Paraná river basin, Reservoirs, Species distribution model

References

- [1] Araujo MB, Guisan A, 2006. Five (or so) challenges for species distribution modelling. *J. Biogeogr.* 33:1677-1688.
- [2] Beaumont LJ, Gallagher RV, Thuiller W, Downey PO, Leishmann MR, Hughes L, 2009. Different climatic envelopes among invasive populations may lead to underestimations of current and future biological invasion. *Divers. Distrib.* 15:409-420.

- [3] Behning AL, 1941. The Cladocerans of the Caucasus. Gruzmedgiz Publishing, Tbilisi: 383 pp.
- [4] Benzie JAH, 2005. Cladocera: The genus *Daphnia* (including *Daphniopsis*) (Anomopoda: Daphniidae). Backhuys Publishers, Leiden: 376 pp.
- [5] Broennimann O, Guisan A, 2008. Predicting current and future biological invasions: both native and invaded ranges matter. *Biol. Lett.* 4:585-589.
- [6] Broennimann O, Treier U.A, Meuller-Schearer H, Thuiller W, Peterson AT, Guisan A, 2007. Evidence of climatic niche shift during biological invasion. *Ecol. Lett.* 10:701-709.
- [7] Brown RS, 1999. Aspects of the biology of *Daphnia lumholtzi* (Sars) in Ft. Loudoun Reservoir, Tennessee. Ph.D. Thesis, University of Tennessee, Knoxville.
- [8] Campbell LP, Luther C, Moo-Llanes D, Ramsey JM, Danis-Lozano R, Peterson AT, 2015. Climate change influences on global distributions of dengue and Chikungunya virus vectors. *Phil. T. R. Soc. B.* 370: 20140135.
- [9] Capinha C, Leung B, Anastacio P, 2011. Predicting worldwide invasiveness for four major problematic decapods: an evaluation of using different calibration sets. *Ecography* 34:448-459.
- [10] Carlton JT, Geller JB, Reaka-Kudla ML, Norse EA, 1999. Historical extinctions in the sea. *Ann. Rev. Ecol. Syst.* 30:515-538.
- [11] Chambers PA, Lacoul P, Murphy KJ, Thomaz SM, 2008. Global diversity of aquatic macrophytes in freshwater. *Hydrobiologia* 595:9-26.
- [12] Clavero M, García-Berthou E, 2005. Invasive species are a leading cause of animal extinctions. *Trends Ecol. Evol.* 20: 110.
- [13] Colautti RI, MacIsaac HJ, 2004. A neutral terminology to define 'invasive' species. *Divers. Distrib.* 10:135-141.
- [14] Colautti RI, Bailey SA, van Overdijk CDA, Admunsen K, MacIsaac HJ, 2006. Characterised and projected costs of nonindigenous species in Canada. *Biol. Invasions* 8:45-59.
- [15] Costa GC, Nogueira C, Machado RB, Colli GR, 2010 Sampling bias and the use of ecological niche modeling in conservation planning: a field evaluation in a biodiversity hotspot. *Bio. Conserv.* 19: 883-899.
- [16] Cox GW, 2004. Alien species and evolution. Island Press, Washington DC: 400 pp.
- [17] De Marco JRP, Siqueira MF, 2009. [Como determinar a distribuição de espécies sobre uma abordagem conservacionista]. [Article in Portuguese]. *Megadiversidade* 5:65-769.
- [18] Dobberfuhl DL, Elser JJ, 2002. Distribution and potential competitive effects of an exotic zooplankter (*Daphnia lumholtzi*) in Arizona reservoirs. *J. Arizona-Nevada Acad. Sci.* 34:89-94.
- [19] Dobrzycka-Kraheil A, Graca B, 2014. Laboratory study of the effect of salinity and ionic composition of water on the mortality and osmoregulation of the gammarid amphipod *Dikerogammarus haemobaphes* (Eichwald, 1841): implications for understanding its invasive distribution pattern. *Mar. Freshwater Res.* 47:227-238.
- [20] Dobrzycka-Kraheil A, Melzer M, Majkowski W, 2015. Range extension of *Dikerogammarus villosus* (Sowinsky, 1894) in Poland (the Baltic Sea basin) and its ability to osmoregulate in different environmental salinities. *Oceanol. Hydrobiol. St.* 44: 94-304.
- [21] Dzialowski AR, Lennon JT, O'Brien WJ, Smith VH, 2003. Predator-induced phenotypic plasticity in the exotic cladoceran *Daphnia lumholtzi*. *Freshwater Biol.* 48:1593-1602.
- [22] Eliás-Gutiérrez M, Martínez-Jerónimo F, Ivanova NV, Valdez-Moreno M, Hebert PDN, 2008. DNA barcodes for Cladocera and Copepoda from Mexico and Guatemala, highlights and new discoveries. *Zootaxa* 1839:1-42.
- [23] Elith J, Graham CH, Anderson RP, Dudik M, Lohmann LG, Loiselle BA, Manion G, Moritz C, Nakamura M, Nakazawa Y, Overton JMcCM, Townsend Peterson A, Phillips SJ, Richardson K, Scachetti-Pereira R, Schapire RE, Soberón J, Williams S, Wisz MS, Zimmermann NE, 2006. Novel methods improve prediction of species' distributions from occurrence data. *Ecography* 29:129-151.
- [24] Engel K, Tollrian R, 2009. Inducible defences as key adaptations for the successful invasion of *Daphnia lumholtzi* in North America? *P. Roy. Soc. Lond. B-Biol.* 276:1865-1873.
- [25] Engel K, Schreder T, Tollrian R, 2014. Morphological defences of invasive *Daphnia lumholtzi* protect against vertebrate and invertebrate predators. *J. Plankton. Res.* 36:1140-1145.
- [26] Fey SB, Cottingham KL, 2012. Thermal sensitivity predicts the establishment success of non-native species in a mesocosm warming experiment. *Ecology* 93:2313-2320.
- [27] Grandcolas P, Nattier R, Trewick S, 2014. Relict species: a relict concept? *Trends Ecol. Evol.* 29:655-663.
- [28] Guisan A, Thuiller W, 2005. Predicting species distribution: offering more than simple habitat models. *Ecol. Lett.* 8:993-1009.
- [29] Guisan A, Baumgartner JB, Naujokaitis-Lewis I, Sutcliffe RP, Tulloch AIT, Regan TJ, Brotons L, McDonald-Madden E, Mantyka-Pringle C, Martin TG, Rhodes JR, Maggini R, Setterfield SA, Elith J, Schwartz MW, Wintle BA, Broennimann O, Austin M, Ferrier S, Kearney MR, Possingham HP, Buckley YM, 2013. Predicting species distributions for conservation decisions. *Ecol. Lett.* 16:1424-1435.
- [30] Hammer Ø, Harper DAT, Ryan PD, 2001. PAST: Paleontological Statistics Software Package for education and data analysis. *Palaeontol. Electron.* 4:1-9.

- [31] Hann BJ, 1986. Revision of the genus *Daphniopsis* Sars, 1903 and description of *Daphniopsis chilensis*, new species from South America. *J. Crustacean Biol.* 6:246-263.
- [32] Havel JE, Hebert PDN, 1993. *Daphnia lumholtzi* in North America: another exotic zooplankter. *Limnol. Oceanogr.* 38:1823-1827.
- [33] Havel JE, Medley KA, 2006. Biological invasions across spatial scales: intercontinental, regional, and local dispersal of cladoceran zooplankton. *Biol. Invasions.* 8:459-473.
- [34] Havel JE, Colbourne JK, Hebert PDN, 2000. Reconstructing the history of intercontinental dispersal in *Daphnia lumholtzi* by use of genetic markers. *Limnol. Oceanogr.* 45: 1414-1419.
- [35] Havel JE, Lee CE, Zanden MJV, 2005. Do reservoirs facilitate invasions into landscapes? *BioScience* 55:518-525.
- [36] Havel JE, Kenko KE, Thomaz SM, Amalfitano S, Kats LB, 2015. Aquatic invasive species: challenges for the future. *Hydrobiologia* 750:147-170.
- [37] Havens KE, East TL, Marcus J, Essex P, Bolan B, Raymond S, et al., 2000. Dynamics of the exotic *Daphnia lumholtzi* and native macro-zooplankton in a subtropical chain-of-lakes in Florida, U.S.A. *Freshwater Biol.* 45:21-32.
- [38] Havens KE, Beaver JR, East TL, Work K, Philips DJ, Cichra MF, Croteau AC, Rodusky AJ, Fulton RS, Rosati TC, 2012. The outcome of the invasion of Florida lakes by *Daphnia lumholtzi*. *Freshwater Biol.* 57: 552-562.
- [39] Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A, 2005. Very high resolution interpolated climate surfaces for global land areas. *Int. J. Climatol.* 25:1965-1978.
- [40] Hulme PE, 2006. Beyond control: wider implications for the management of biological invasions. *J. Appl. Ecol.* 43:835-847.
- [41] Incagnone G, Marrone F, Barone R, Robba L, Naselli-Flores L, 2015. How do freshwater organisms cross the "dry ocean"? A review on passive dispersal and colonization processes with a special focus on temporary ponds. *Hydrobiologia* 750:103-123.
- [42] Jiménez-Valverde A, Lobo JM, Hortal J, 2008. Not as good as they seem: the importance of concepts in species distribution modelling. *Divers. Distrib.* 14:885-890.
- [43] Jimenez-Valverde A, Peterson AT, Sobéron J, Overton JM, Aragon P, Lobo JM, 2011. Use of niche models in invasive species risk assessments. *Biol. Invasions* 13:2785-2797.
- [44] Johnson JL, Havel JE, 2001. Competition between native and exotic *Daphnia*: in situ experiments. *J. Plankton Res.* 23: 373-387.
- [45] Johnson PTJ, Olden JD, Vander Zanden MJ, 2008. Dam invaders: impoundments facilitate biological invasions into freshwaters. *Front. Ecol. Environ.* 6:357-363.
- [46] Kolar CS, Boase JC, Clapp DF, Whal DH, 1997. Potential effect of invasion by an exotic zooplankter, *Daphnia lumholtzi*. *J. Freshwater Ecol.* 12:521-530.
- [47] Kotov AA, Taylor DJ, 2010. A new African lineage of the *Daphnia obtusa* group (Cladocera: Daphniidae) disrupts continental vicariance patterns. *J. Plankton Res.* 32:937-949.
- [48] Kotov AA, Taylor DJ, 2014. *Daphnia lumholtzi* Sars, 1885 (Cladocera: Daphniidae) invades Argentina. *J. Limnol.* 73:364-374.
- [49] Kotov AA, Sinev AY, Berrios VL, 2010. The Cladocera (Crustacea: Branchiopoda) of six high altitude water bodies in the North Chilean Andes, with discussion of Andean endemism. *Zootaxa* 2430:1-66.
- [50] Kotov AA, Forró L, Korovchinsky NM, Petrusek A, 2013. World checklist of freshwater Cladocera species. World Wide Web electronic publication. Available from: <http://fada.biodiversity.be/group/show/17>
- [51] Lavergne S, Molina J, Debussche MAX, 2006. Fingerprints of environmental change on the rare mediterranean flora: a 115-year study. *Glob. Change Biol.* 12: 466-1478.
- [52] Lenon JT, Smith VH, Williams K, 2001. Influence of temperature on exotic *Daphnia lumholtzi* and implications for invasion success. *J. Plankton Res.* 23:25-434.
- [53] Marcatonio M, Mets M, Baldachino F, Arnoldi D, Montarsi F, Campelli G, Carlin S, Neteler M, Rizzoli A, 2016. First assessment of potential distribution and dispersal capacity of the emerging invasive mosquito *Aedes koreicus* in Northeast Italy. *Parasit. Vectors.* 9: 63.
- [54] McMahon RF, 2002. Evolutionary and physiological adaptation of aquatic invasive animal: r selection versus resistance. *Can. J. Fish. Aquat. Sci.* 59:1235-1244.
- [55] Mergeay J, Verschuren D, De Meester L, 2005a. Cryptic invasion and dispersal of an American *Daphnia* in East Africa. *Limnol. Oceanogr.* 50: 1278-1283.
- [56] Mergeay J, Verschuren D, De Meester L, 2005b. *Daphnia* species diversity in Kenya, and a key to the identification of their ephippia. *Hydrobiologia.* 542: 261-274.
- [57] Mergeay J, Verschuren D, De Meester L, 2006. Invasion of an asexual American water flea clone throughout Africa and rapid displacement of a native sibling species. *P. Roy. Soc. Lond. B-Biol.* 273:2839-2844.
- [58] Mooney HA, Cleland EE, 2001. The evolutionary impact of invasive species. *Proc. Nat. Acad. Sci. USA.* 98: 5446-5451.

- [59] Nori J, Urbina-Cardona JN, Loyola RD, Lescano JN, Leynaud GC, 2011. Climate change and American bullfrog invasion: what could we expect in South America? PLoS One 6:e25718.
- [60] Padhye SM, Kotov AK, Dahanukar N, Dumont HJ, 2016. Biogeography of *Daphnia* O. F. Müller (Crustacea: Branchiopoda: Anomopoda) on the Indian subcontinent. J. Limnol. 75:571-580.
- [61] Palaoro AV, Dalosto MM, Costa GC, Santos S, 2013. Niche conservatism and the potential for the crayfish *Procambarus clarkii* to invade South America. Freshwater Biol. 58:1379-1391.
- [62] Panov VE, Krylov PI, Riccardi N, 2004. Role of diapause in dispersal and invasion success by aquatic invertebrates. J. Limnol. 63:56-69.
- [63] Perrings C, Dehnen-Schmutz K, Touza J, Williamson M, 2005. How to manage biological invasions under globalization. Trends Ecol. Evol. 20:212-215.
- [64] Peterson AT, 2003. Predicting the geography of species' invasions via ecological niche modeling. Q. Rev. Biol. 78:419-433.
- [65] Petitpierre B, Kueffer C, Broennimann O, Randin C, Daehler C, Guisan A, 2012. Climatic niche shifts are rare among terrestrial plant invaders. Science 335:1344-1348.
- [66] Phillips SJ, Anderson RP, Schapire RE, 2006. Maximum entropy modeling of species geographic distributions. Ecol. Modell. 190:231-259.
- [67] Pitman NC, Jørgensen PM, Williams RS, León-Yáñez S, Valencia R, 2002. Extinction-Rate Estimates for a Modern Neotropical Flora. Conser. Biol. 16:1427-1431.
- [68] Poulin R, Paterson RA, Townsend CR, Tompkins DM, Kelly DW, 2011. Biological invasions and the dynamics of endemic diseases in freshwater ecosystems. Freshwater Biol. 56:676-688.
- [69] Rodríguez JP, 2001. Exotic species introductions into South America: an underestimated threat? Biod. Conserv. 10:1983-1996.
- [70] Silva-Briano M, Arroyo-Bustos G, Beltran-Alvarez R, Adabache-Ortiz A, Galvan de la Rosa R, 2010. [*Daphnia Ctenodaphnia lumholtzi* G.O. Sars, 1885 (Crustacea: Cladocera); an exotic cladoceran in Mexico]. [Article in Spanish with English Abstract]. Hidrobiológica 20:275-280.
- [71] Simberloff D, 2010. Invasive species, p. 131-152. In: N.S. Sodhi and P.R. Ehrlich (eds.), Conservation biology for all. Oxford Biology.
- [72] Simões NR, Robertson BA, Lansac-Tôha FA, Takahashi EM, Bonecker CC, Velho LFM, Joko CY, 2009. Exotic species of zooplankton in the Upper Paraná River floodplain, *Daphnia lumholtzi* Sars, 1885 (Crustacea: Branchiopoda). Braz. J. Biol. 69:551-558.
- [73] Soberon J, 2007. Grinnellian and Eltonian niches and geographic distribution of species. Ecol. Lett. 10:1115-1123.
- [74] Sodhi NS, Brook BW, Bradshaw CAJ, 2009. Causes and consequences of species extinctions, p. 514-520. In: S.A. Leviton (ed.), Princeton guide to ecology. Princeton University Press.
- [75] Sousa FDR, Elmoor-Loureiro LMA, Palaoro AV, Silvano DL, 2014. [Predição da invasão de *Daphnia lumholtzi* (Crustacea, Branchiopoda) na região Neotropical]. [Article in Portuguese]. Proceedings 30th Congresso Brasileiro de Zoologia, Porto Alegre.
- [76] Speziale K, Lambertucci S, 2010. A call for action to curb invasive species in South America. Nature 467:153-153.
- [77] Valdivia-Vilar RS, Burger L, 1989. Description of *Daphniopsis marcahuasensis* sp. nov. (Cladocera: Daphniidae) from Peru with the inclusion of a key of the genus species. Amazoniana 10:439-452.
- [78] Wittmann MJ, Gabriel W, Harz EM, Laforsch C, Jeschke JM, 2013. Can *Daphnia lumholtzi* invade European lakes? Neo-Biota 16:39-57.
- [79] Work KA, Gophen M, 1999. Environmental variability and the population dynamics of the exotic *Daphnia lumholtzi* and native zooplankton in Lake Texoma, U.S.A. Hydrobiologia 405:11-23.
- [80] Zanata LH, Espindola ELG, Rocha O, Pereira RHG, 2003. First record of *Daphnia lumholtzi* (Sars, 1885), exotic cladoceran, in São Paulo State (Brazil). Braz. J. Biol. 63:717-720.
- [81] Zanata LH, Espindola ELG, Rocha O, Pereira RHG, 2008. Morphological abnormalities in Cladocera (Branchiopoda) in a cascade of reservoirs in the middle and lower Tietê river (São Paulo, Brazil). Braz. J. Biol. 68:681-682.
- [82] Zhang YH, Wang JJ, Comes HP, Peng H, Qiu YX, 2016. Contributions of historical and contemporary geographic and environmental factors to phylogeographic structure in a Tertiary relict species, *Emmenopterys henryi* (Rubiaceae). Sci. Rep. 6: 24041.